

END NOTES

Foreword

Page xi: “The incidence of childhood cancer is up about 25%...” and “more than six million children now suffer from asthma” - U.S. Environmental Protection Agency, 2003. America’s Children and the Environment: Measures of Contaminants, Body Burdens, and Illnesses. http://www.epa.gov/economics/children/publications/ace_2003.pdf

Page xi: “One in every thirty-three babies born in the United States enters the world with a birth defect.” - U.S. Centers for Disease Control and Prevention, 2007. National Center on Birth Defects and Developmental Disabilities. <http://www.cdc.gov/ncbddd/bd/faq1.htm#chanceofBD>

Page xi: “One in six children deals with at least one developmental disorder.” - U.S. Centers for Disease Control and Prevention, 2007. National Center on Birth Defects and Developmental Disabilities. <http://www.cdc.gov/ncbddd/dd/ddsurv.htm>

Page xi: “Between 3-5 percent of children have Attention Deficit Hyperactivity Disorder.” – National Institute of Mental Health, 2006. Attention Deficit Hyperactivity Disorder. <http://www.nimh.nih.gov/health/publications/adhd/nimhadhdpub.pdf>

Introduction

Page 2: “42 billion – Number of pounds of chemicals the United States produces or imports each day.” - National Pollution Prevention and Toxics Advisory Committee (NPPTAC), 2005. Broader Issues Work Group. How can EPA more efficiently identify potential risks and facilitate risk reduction decision for non-HPV existing chemicals? <http://www.epa.gov/oppt/npptac/pubs/finaldraftnonhpvpaper051006.pdf>

Page 4: “The incidence of childhood cancers jumped almost 27% between 1975 and 2002” - U.S. Environmental Protection Agency, 2003. America’s Children and the Environment: Measures of Contaminants, Body Burdens, and Illnesses. http://www.epa.gov/economics/children/publications/ace_2003.pdf

Page 4: “Treatments kill cancer cells, but they damage healthy cells as well, causing complications from heart problems and liver failure to infertility.”

National Cancer Institute, 2007. Late Effects of Treatment for Childhood Cancer. <http://www.cancer.gov/cancertopics/pdq/treatment/lateeffects/Patient/page1>

American Cancer Society, 2007. Childhood Cancer: Late Effects of Cancer Treatment. http://www.cancer.org/docroot/CRI/content/CRI_2_6x_Late_Effects_of_Childhood_Cancer.asp

Page 4: “Survivors of childhood cancer have a mortality rate more than ten times higher than the general population’s due to the long term effects of treatment.” Journal of Clinical Oncology, 2001 (Vol 19, Issue 13). Late Mortality Experience in Five-Year Survivors of Childhood and Adolescent Cancer: The Childhood Cancer Survivor Study. <http://jco.ascopubs.org/cgi/content/full/19/13/3163>

Page 4: “The American Cancer Society estimates that 75 percent of cancer is due to environmental factors.” – American Cancer Society, 2007. What Are the Risk Factors for Cancer? (environmental factors include tobacco use, diet, infectious disease, chemicals and radiation). http://www.cancer.org/docroot/CRI/content/CRI_2_4_2x_What_are_the_risk_factors_for_cancer_72.asp

Page 5: “According to the US Centers for Disease Control and Prevention, in 2005 6.5 million children under age eighteen had asthma; an increase of more than 200% since 1980.” US Centers for Disease Control, National Center for Health Statistics, 2006. State of Childhood Asthma, United States: 1980-2005. <http://www.cdc.gov/nchs/data/ad/ad381.pdf>

Page 5: “About one in eleven school-aged children suffers from asthma, and the rate is rising more rapidly in preschool-aged children than in any other age group.” US EPA, 2007. Indoor Air Quality. <http://www.epa.gov/iaq/>

Page 5: “Indoor air quality is a big culprit. In addition to dust mites, mold, pet dander, and secondhand smoke, air contaminants that may impact asthma include certain insecticides and chemicals in plastic, especially formaldehyde.” US EPA, 2007. Indoor Air Quality. <http://www.epa.gov/iaq/> European Respiratory Journal, 2002 (Vol 20, Issue 2). Domestic Exposure to Formaldehyde Significantly Increases the Risk of Asthma in Young Children. <http://www.erj.ersjournals.com/cgi/content/full/20/2/403>

Page 5: “Allergic dermatitis (itchy rash) is the most common skin condition in children younger than 11 years of age and the percentage of children diagnosed with it has increased over 300% from 3 percent in the 1960s to 10 percent in the 1990s. “ – American Academy of Allergy Asthma & Immunology, 2007. Patient/Public Education: Fast Facts. http://www.aaaai.org/patients/resources/fastfacts/skin_conditions.stm

Page 5: “Hay fever (allergic rhinitis) is believed to affect up to 40% of children.” Management of Allergic and Nonallergic Rhinitis. Summary, Evidence Report/Technology Assessment: Number 54, May, 2002. Rockville, Md.: U.S. Department of Health and Human Services, Agency for Healthcare Research and Quality. <http://www.ncbi.nlm.nih.gov/books/bv.fcgi?rid=hstat1.chapter.117840>

Page 5: “On any given day, ten thousand American children miss school because of hay fever, for an annual total of 2 million lost school days.” American Academy of Allergy, Asthma & Immunology, 2007. Allergy Statistics. http://www.aaaai.org/media/resources/media_kit/allergy_statistics.stm

Page 5: “roughly six percent of children suffer from food allergies” US CDC. Allergic conditions among children: US, 2000-2005 (Source: NHIS) Available online at <http://209.217.72.34/HDAA/TableViewer/tableView.aspx?ReportId=205>. Accessed September 17, 2007.

Report on the Expert Panel on Food Allergy Research, June 30 and July 1, 2003, National Institute of Allergy and Infectious Diseases, National Institutes of Health. http://www3.niaid.nih.gov/about/organization/dait/PDF/june30_2003.pdf [PDF 190K].

Page 5: “Diagnoses for ADHD has jumped almost 400%” - Mandell DS, Thompson WW, Weintraub ES, et al. Trends in the diagnosis rates for autism and ADHD at hospital discharge in the context of other psychiatric diagnoses. *Psychiatr Serv.* 2005;56:56-62. <http://www.medscape.com/viewarticle/511173>

Page 5: “estimates that up to 600,000 children born each year in the United States suffer mild to moderate loss of IQ points because of methylmercury exposure during fetal development.” American Public Health Association, 2008. <http://www.apha.org/about/news/pressreleases/2008/CAMR.htm>

Page 6: “Endocrine-disruptors—often found in plastics and cosmetics - mimic or block hormones, disrupting normal function, and have been shown to cause birth defects in laboratory animals.” Environmental Health Perspectives, 2003 (October). Assessing the Effects of Endocrine Disruptors in the National Children’s Study. <http://www.ehponline.org/members/2003/5799/5799.html#gaps>

Page 6: “One in six American children between the ages of six and 19 are considered overweight.” Epidemiologic Reviews, 2007 (May). The Obesity Epidemic in the United States – Gender, Age, Socioeconomic, Racial/Ethnic, and Geographic Characteristics: A Systematic Review and Meta-Regression Analysis <http://epirev.oxfordjournals.org/cgi/content/abstract/29/1/6>

Page 6: “Type 2 diabetes, once known as adult-onset diabetes, is increasing among children. One in three children born in the year 2000 will develop diabetes; for an African-American or Hispanic baby, the ratio is closer to 1 in 2.” US Centers for Disease Control, 2005. Preventing Diabetes and Its Complications. <http://www.cdc.gov/nccdphp/publications/factsheets/Prevention/diabetes.htm>

Page 6: “Studies have shown that chemicals, like Bisphenol-A (BPA)...may interfere with normal hormone function, making obesity more likely.” Journal of Lipid Research, 2002 (May). Bisphenol A in Combination with Insulin Can Accelerate the Conversion of 3T3-L1 Fibroblasts to Adipocytes. <http://www.jlr.org/cgi/content/abstract/43/5/676>

Toxicological Sciences, 2003. Endocrine Disruptors and the Obesity Epidemic. <http://toxsci.oxfordjournals.org/cgi/content/full/76/2/247?ck=nck>

Chapter 1: Doing the Bump: Preparing for Baby

Page 15: “2005 study by the Environmental Working Group that examined cord blood from ten babies born during a two-month span in US hospitals. What they found is heart-stopping: Present in the cord blood was an average of 200 chemicals and pollutants commonly found in the home, including mercury, flame retardants, perfluorinated chemicals (from nonstick surfaces and carpet), plasticizers (PVCs), pesticides, and wood preservatives—many of which may cause cancer, are toxic to the brain and nervous system, and are linked to birth defects or abnormal development” Environmental Working Group, 2005. *Body Burden – The Pollution in Newborns*.
<http://archive.ewg.org/reports/bodyburden2/execsumm.php>

Page 16: “*You can lower your pesticide exposure by 90% simply by avoiding the dozen most contaminated fruits and vegetables, identified by the Environmental Working Group.*” Environmental Working Group, 2007. *In the Ideal Position to Avoid Pesticides*.
<http://www.ewg.org/node/21381>

Page 16: “some experts have suggested that a high consumption of the hormonally active natural plant substances (phytoestrogens) present in soy explain the low incidence of estrogen-related cancers (like breast cancer) and heart disease in Asian women.” Environmental Health Perspectives, 1995 (October). *Phytoestrogens: Epidemiology and a Possible Role in Cancer Protection*. [http://links.jstor.org/sici?sici=0091-6765\(199510\)103%3C103%3APEAAPR%3E2.0.CO%3B2-H](http://links.jstor.org/sici?sici=0091-6765(199510)103%3C103%3APEAAPR%3E2.0.CO%3B2-H)

The Journal of Clinical Endocrinology & Metabolism, 1998. *Potential Health Benefits of Dietary Phytoestrogens: A Review of the Clinical, Epidemiological, and Mechanistic Evidence*.
http://jcem.endojournals.org/cgi/content/full/83/7/2223?maxtoshow=&HITS=40&hits=40&RESULTFORMAT=1&titleabstract=phytoestrogen&fulltext=soy&searchid=1056834341692_1156&stored_search=&FIRSTINDEX=0&sortspec=relevance&tdate=6/30/2003&resourcetype=1

Page 16: “studies suggest that overexposure to phytoestrogens can be harmful to developing babies” Naz, Rajesh, ed. 2004. *Endocrine Disruptors: Effects on Male and Female Reproductive Systems*, Second Edition. CRC Press.

US EPA, 1999. *Developmental Effects of Dietary Soy Phytoestrogens*.
http://es.epa.gov/ncer/science/endocrine/pdf/development/r825721_hughes_0415_draft.pdf

Page 17: “From now on try to pick up organic meat and milk, minimizing your exposure to the antibiotics and hormones used in their production, all of which could be bad for mom’s health—ergo, bad for baby’s.” Health Care Without Harm. *Position Statement on rBGH*. <http://www.noharm.org/details.cfm?ID=1104&type=document>

The Scientific Committee on Veterinary Measures Relating to Public Health.
“Assessment of Potential Risks to Human Health from Hormone Residues in Bovine Meat and Meat Products.” European Commission, April 30, 1999.

Page 17: “large species, including swordfish, king mackerel and, sorry to say, that sandwich-staple, tuna, may contain elevated levels of mercury, which can upset normal fetal brain growth. Chunk light tuna is preferable to white albacore, because it typically has lower levels of mercury (it comes from smaller fish). The EPA advises that pregnant women can safely have up to 12 ounces of chunk light tuna a week.” US Environmental Protection Agency, 2007. What You Need to Know About Mercury in Fish and Shellfish. <http://www.epa.gov/waterscience/fishadvice/advice.html>

Page 17: “The way they’re raised means farmed fish often contain in their fat significant concentrations of contaminants, like PCBs—a banned synthetic substance linked to developmental delays in children exposed pre-natally (and which can linger in human fat for decades).” Chemosphere, 2007. Chemical Contaminants in Fish Feeds Used in Federal Salmonid Hatcheries in the USA. http://www.sciencedirect.com/science?_ob=ArticleURL&_udi=B6V74-4MT552N-1&_user=10&_rdoc=1&_fmt=&_orig=search&_sort=d&_view=c&_acct=C000050221&_version=1&_urlVersion=0&_userid=10&md5=1241bbf2aa88b6e12963ba28e73a0417

Page 17: “600,000 - Estimated number of babies born each year with a mild to moderate loss of IQ, thanks to in-utero exposure to methylmercury. Fish consumption is the likely culprit.” US Environmental Protection Agency, 2004. Referenced by the American Public Health Association. <http://www.apha.org/about/news/pressreleases/2008/CAMR.htm>

Page 17: “*Listeria* is one: The CDC claims that pregnant women are 20 times more likely to become infected than non-pregnant adults.” US Centers for Disease Control, 2005. Division of Bacterial and Mycotic Diseases. http://www.cdc.gov/ncidod/dbmd/diseaseinfo/listeriosis_g.htm

Page 18: “If mom-to-be does get sick, the consequences for baby could be grave, from miscarriage to infection in the newborn.” World Health Organization, 2007. *Listeria* Infections. http://www.who.int/topics/listeria_infections/en/

Page 18: “The CDC recommends that pregnant women avoid soft cheeses such as feta, brie, camembert, and blue-veined cheeses.” US Centers for Disease Control, 2005. Division of Bacterial and Mycotic Diseases. http://www.cdc.gov/ncidod/dbmd/diseaseinfo/listeriosis_g.htm

Page 18: “Toxoplasmosis, an infection from a parasite in cat feces, can threaten the health of an unborn child.” American Journal of Medicine, 2005. Toxoplasmosis in Pregnancy. <http://www.ncbi.nlm.nih.gov/sites/entrez?db=pubmed&uid=15745715&cmd=showdetailview&indexed=google>

Page 19: “Monosodium glutamate is a flavor enhancer found in Asian cuisine and soups that can cause headaches and stomach upset.” US Department of Health and Human Services, 1995. FDA and Monosodium Glutamate (MSG).

<http://www.cfsan.fda.gov/~lrd/msg.html>

Page 19: “At high concentrations, it’s also a neurotoxin (kills nerve cells).” Endocrinology, 1984. Increased responsiveness of the hypothalamic-pituitary axis after neurotoxin-induced hypothalamic denervation.

<http://www.ncbi.nlm.nih.gov/sites/entrez?db=pubmed&uid=6145582&cmd=showdetailview&indexed=google>

Page 19: “steer clear of are blue 1 & 2, green 3, red 3, and yellow 6.” Townsend Letter, 2007. Artificial Food and Cosmetic Coloring: Hidden Sources of Toxic Metals.

<http://www.townsendletter.com/Nov2007/artfood1107.htm>

Page 19: “aspartame has been linked to cancer in animal studies” Environmental Health Perspectives, 2006. First experimental demonstration of the multipotential carcinogenic effects of aspartame administered in the feed to Sprague-Dawley rats.

http://www.ncbi.nlm.nih.gov/sites/entrez?db=PubMed&cmd=Retrieve&list_uids=16507461

Page 19: “Nitrites. Food additives normally found in hot dogs and sandwich meats, these are possibly carcinogenic.” Journal of the National Cancer Institute, 2005. Meat and Fat Intake as Risk Factors for Pancreatic Cancer: The Multiethnic Cohort Study.

<http://jnci.oxfordjournals.org/cgi/content/abstract/97/19/1458>

Page 20: “A recent study from Tufts University revealed some worrisome results from a very commonly used chemical called Bisphenol-A (BPA), contained in clear bicarbonate plastics often used for water bottles and baby bottles. BPA can mimic estrogen and mess with reproductive growth.” Endocrinology, 2005. Perinatal Exposure to Bisphenol-A Alters Peripubertal Mammary Gland Development in Mice.

<http://www.tufts.edu/sackler/cmdb/soto-lab/publications/pdf/11.pdf>

Molecular and Cellular Endocrinology, 2006. Endocrine Disruptors and Reproductive Health: The Case of Bisphenol-A. <http://www.tufts.edu/sackler/cmdb/soto-lab/publications/pdf/5.pdf>

Page 20: “Also, bottled water is less regulated than tap so there's no guarantee it's any safer.” Natural Resources Defense Council, 1999. Bottled Water: Pure Drink or Pure Hype? <http://www.nrdc.org/water/drinking/bw/exesum.asp>

The New York Times, 2005. Bad to the Last Drop.

<http://www.elporvenir.org/article01.html>

Page 23: “If you use prescription skin medications, such as Accutane for acne or topical steroids for eczema, you’ll want to give them up during pregnancy: No less than the CDC

advises it” US Centers for Disease Control, 2000. Accutane Exposed Pregnancies. <http://www.cdc.gov/mmwr/preview/mmwrhtml/mm4902a2.htm>

Page 23: “Some hair color products may contain harmful chemicals such as lead acetate and coal tar dyes.” US Food and Drug Administration, 1997. Hair Dye Products. <http://www.cfsan.fda.gov/~dms/cos-hdye.html>

Page 23: “Indoor air—the stuff we sleep in, play in, eat in; the air you might dare call your own—can be far more polluted than the air outside; in fact, it’s as much as *two to five times worse*.” US EPA, 2004. Buildings and the Environment: A Statistical Summary. <http://www.epa.gov/opptintr/greenbuilding/pubs/gbstats.pdf>.

Page 24: “Toxic chemicals including lead and pesticides are routinely found in household dust.” Ecology Center, 2005. Sick of Dust: Chemicals in Common Products – A Needless Health Risk in Our Homes. http://ecocenter.org/releases/dust_rpt/Dust-Report.pdf

Environmental Working Group, 2004. In the Dust: Toxic Fire Retardants in American Homes. http://www.ewg.org/files/InTheDust_final.pdf

Page 24: “In one study (funded by Healthy Child), researchers at the Columbia University Center for Children’s Environmental Health evaluated 60 newborns whose mothers wore portable air monitors during their last trimester and found that babies’ DNA can undergo subtle damage by the polluted air their mothers breathe during pregnancy.” Cancer Epidemiology, Biomarkers & Prevention, 2005. Chromosomal Aberrations in Cord Blood Are Associated with Prenatal Exposure to Carcinogenic Polycyclic Aromatic Hydrocarbons. <http://cebp.aacrjournals.org/cgi/content/full/14/2/506>

Page 24: “Tobacco smoke components, including nicotine, can quickly cross the placenta and lead to birth defects...” Pediatrics, 2004. Prenatal and Postnatal Environmental Tobacco Smoke Exposure and Children’s Health. <http://pediatrics.aappublications.org/cgi/content/full/113/4/S1/1007>

Page 24: “...mothers who smoke or are exposed to secondhand smoke during pregnancy are more likely to give birth to low-birth weight babies.” Epidemiology, 2000. Prenatal Active or Passive Tobacco Smoke Exposure and the Risk of Preterm Delivery or Low Birth Weight. <http://www.epidem.com/pt/re/epidemiology/abstract.00001648-200007000-00011.htm;jsessionid=HIPLpbR6HLp1Rg9JnqzRKpLsyS3yQTyyP7TT7QlgvvNS1dw1mxFf!1219373867!181195629!8091!-1>

Page 24: “But conventional cleaning products contain toxic chemicals such as ammonia, formaldehyde, and artificial fragrances that may pose a risk if a pregnant woman is exposed. One study from the University of Bristol in the UK found that babies of moms who frequently used chemical-based cleaners while pregnant were more than twice as likely to develop breathing problems.” Sherriff A, Farrow A, Golding J, ALSPAC Study Team, Henderson AJ. Frequent use of chemical household products is associated with

persistent wheezing in preschool-age children. *Thorax* 2005; 60: 45-9.

Page 25: “Commercial air fresheners contain harmful volatile chemicals.” National Institutes of Health, 2006. Chemicals in Many Air Fresheners May Reduce Lung Function. <http://www.nih.gov/news/pr/jul2006/nihs-27.htm>

Natural Resources Defense Council, 2007. Protect Your Family from the Hidden Hazards in Air Fresheners. <http://www.nrdc.org/health/home/airfresheners/fairfresheners.pdf>

Page 25: “Even scented candles are a dubious option, as they can release harmful carcinogens and some may even contain lead.” US EPA, 2001. Candles and Incense as Potential Sources of Indoor Air Pollution: Market Analysis and Literature Review.

Page 28: “Since the VOC fumes that waft from conventional paint (in part from the solvents that keep paints wet) may be harmful to your baby and can linger for weeks,” US Centers for Disease Control, 2005. Having a Healthy Pregnancy. <http://www.cdc.gov/ncbddd/bd/abc.htm>

Page 28: “Lead poisoning can harm a child’s brain growth, and fetuses can be exposed in utero.” US Centers for Disease Control, 2006. Lead Toxicity: What are the Physiologic Effects of Lead Exposure? http://www.atsdr.cdc.gov/csem/lead/pbphysiologic_effects2.html

Page 31: “Carpet, especially the newly installed stuff, can emit harmful chemicals from fibers, backing material, dyes, and flame-retardants” American Lung Association, 2000. Your Carpet and Indoor Air Quality. <http://www.lungusa.org/site/pp.asp?c=dvLUK9O0E&b=35441>

Page 32: “fiberboard and pressed wood contain a mess of chemical finishes and glues.” US EPA, 2007. Basic Information: Formaldehyde. <http://www.epa.gov/iaq/formalde.html>

Chapter 2 – Clean Up Time: Safe Suds and a Greener Clean

Page 36: “But labels won’t give you the full picture—of the known respiratory irritants, carcinogens, hormone disruptors, and neurotoxins that are associated with chronic and long-term effects—“ California Environmental Protection Agency, 2006. Indoor Air Chemistry: Cleaning Agents, Ozone and Toxic Air Contaminants. <http://www.arb.ca.gov/research/abstracts/01-336.htm#Disclaimer>

Women’s Voices for the Earth, 2007. Household Hazards: Potential Hazards of Home Cleaning Products. <http://www.womenandenvironment.org/campaignsandprograms/SafeCleaning/HazardsReport.pdf>

Page 36: “(No wonder studies show that use of conventional household cleaning products is associated with chronic wheezing and asthma in children.)” *Thorax* (International

Journal of Respiratory Medicine), 2005. Frequent use of chemical household products is associated with persistent wheezing in pre-school age children.
<http://thorax.bmj.com/cgi/content/abstract/60/1/45>

Thorax, 2004. Association of domestic exposure to volatile organic compounds with asthma in young children. <http://thorax.bmj.com/cgi/content/abstract/59/9/746>

Page 36: “To top it off, when these chemicals are washed down the drain, any pretense to being agents of “clean” disappears, as they can pollute our rivers and lakes, poisoning fish and wildlife.” U.S. Geological Survey (2002). Emerging Contaminants Project.
<http://toxics.usgs.gov/regional/emc/index.html>

The Green Guide, 2007. Product Report: Laundry Supplies.
<http://www.thegreenguide.com/reports/productprint.mhtml?id=78>

Ecotoxicology and Environmental Safety, 1999. Toxicity of Laundry Detergent Components to a Freshwater Cladoceran and Their Contribution to Detergent Toxicity.
http://www.sciencedirect.com/science?_ob=ArticleURL&_udi=B6WDM-45GWF8H-V&_user=10&_rdoc=1&_fmt=&_orig=search&_sort=d&_view=c&_acct=C000050221&_version=1&_urlVersion=0&_userid=10&md5=161147c4f0e32aca85993704ddfa8ff9

Page 36: “63 – The number of hazardous chemical products in the average American household, equal to roughly 10 gallons of household hazardous waste” Consumer Product Safety Commission, 2004. Household Hazardous Waste.
http://www.smgov.net/epd/printed_materials/pdf/HHW_Booklet_WEB2004.pdf

Page 37: “Ten percent of the calls to the US Poison Control Centers involve toxic exposure to household cleaners; of these, two-thirds concern children.” American Association of Poison Control Centers, 2005. 2005 Annual Report of the American Association of Poison Control Centers’ National Poisoning and Exposure Database.
<http://www.aapcc.org/Annual%20Reports/05report/2005%20Published.pdf>

Page 38: “*ammonia*, a staple in bathrooms and kitchens for its ability to cut through dirt and grime and kill everything on contact, it’s also a VOC (volatile organic compound), which is carried through the air to your nose as vapor, contributing to respiratory issues, perhaps worse.” US Agency for Toxic Substances & Disease Registry, 2004. Toxicological Profile for Ammonia. <http://www.atsdr.cdc.gov/toxprofiles/tp126.html>

Page 38: “*chlorine*, a component of many bleaching products which can turn into highly hazardous chemicals when interacting with other substances;” US Agency for Toxic Substances & Disease Registry, 2007. Toxicological Profile for Chlorine.
<http://www.atsdr.cdc.gov/toxprofiles/tp172.html#bookmark05>

Page 38: “the #1 household chemical involved in poisoning.” The Green Guide, 2003.

The Chlorine Conundrum. <http://www.thegreenguide.com/doc/95/bleach>

Page 38: “*phosphates*, water-softeners banned from laundry detergents in most states for their damaging effect on fish and freshwater systems; still found in some dishwasher and cleaning formulas.” TreeHugger, 2007. Ask TreeHugger: What’s the Dirt on Phosphate-Free Soaps?

http://www.treehugger.com/files/2007/03/ask_treehugger_whats_the_dirt_on_phosphate-free_soaps.php

Page 39: “*lye* (sodium hydroxide), used in drain and oven cleaners, detergent, pool-cleaners, metal polishers, and soap, it’s made from ash and great at fighting grease and muck; unfortunately, it can also irritate skin and eyes, and lye fumes can corrode respiratory passages.” US Agency for Toxic Substances & Disease Registry, 2002. ToxFAQs for Sodium Hydroxide. <http://www.atsdr.cdc.gov/tfacts178.html>

Page 39: “Synthetic fragrances, which are cocktails of up to hundreds of chemicals, are common triggers of allergic reactions and respiratory distress.” Archives of Environmental Health, 1998. Acute Toxic Effects of Fragrance Products. <http://www.ncbi.nlm.nih.gov/sites/entrez?db=pubmed&uid=9577937&cmd=showdetailview&indexed=google>

Pitts, C. 2003. Get A Whiff Of This: Perfumes – The Invisible Chemical Poisons.

Page 39: “Not all VOCs are harmful, but some have been linked to everything from neurological and organ damage to cancer.” Indoor Air, 1992. Effects of Indoor Air Pollution on Human Health. <http://www.blackwell-synergy.com/doi/abs/10.1111/j.1600-0668.1992.02-21.x>

Page 39: “Recent Australian studies correlated VOC exposure in the home with a higher incidence of asthma.” Thorax (International Journal of Respiratory Medicine), 2005. Frequent use of chemical household products is associated with persistent wheezing in pre-school age children. <http://thorax.bmj.com/cgi/content/abstract/60/1/45>

Thorax, 2004. Association of domestic exposure to volatile organic compounds with asthma in young children. <http://thorax.bmj.com/cgi/content/abstract/59/9/746>

Page 47: “For one, some bacteria are beneficial, as a natural part of our body’s ecosystem, so killing them can damage skin and weaken the immune system.” Journal for Specialists in Pediatric Nursing, 2005. Bactericidal Household Products and Antibiotic Resistance: Issues for Parents.

Page 47: “For another thing, this constant assault of antibacterial products is thought to contribute to the breeding of “super-germs,” or bacteria resistant to disinfectants and drugs” The Lancet Infectious Diseases, 2003. Antibacterial Cleaning and Hygiene Products as an Emerging Risk Factor for Antibiotic Resistance in the Community.

Page 47: “*Triclosan* is the main ingredient in most anti-bacterial and disinfectant soaps and sprays; it’s related to one of our deadliest pollutants, dioxin, which can form when triclosan is mixed with water and exposed to sunlight.” Beyond Pesticides, 2004. The Ubiquitous Triclosan: A Common Antibacterial Agent Exposed.
<http://www.beyondpesticides.org/pesticides/factsheets/Triclosan%20cited.pdf>

Page 48: “66% of streams sampled that were found to contain disinfectants” USGS, 2002. Pharmaceuticals, Hormones, and Other Organic Wastewater Contaminants in US Streams. <http://toxics.usgs.gov/pubs/FS-027-02/pdf/FS-027-02.pdf>

Page 49: “The physical act of washing with soap and water removes all kinds of microorganisms, including the viruses that cause colds and the flu (which, cruel irony, aren’t even destroyed by antibacterials).” Clinical Infectious Diseases, 2007. Consumer Antibacterial Soaps: Effective or Just Risky?
<http://deepblue.lib.umich.edu/dspace/bitstream/2027.42/55477/1/Consumer%20antibacterial%20soaps%20Effective%20or%20just%20risky.web%202007%20aiello.pdf>

Page 49: “275 The number of active ingredients in anti-bacterial products classified as pesticides.” US EPA, 2007. EPA Did Not Properly Process a Hospital Disinfectant and Sanitizer Registration. <http://www.epa.gov/oig/reports/2007/20070329-2007-P-00018.pdf>

Page 49: “A person who spends just 15 minutes cleaning scale off of shower walls could inhale three times the “acute one-hour exposure limit” for glycol ether-containing products set by the California Office of Environmental Health Hazard Assessment.” California Environmental Protection Agency, 2006. Indoor Air Chemistry: Cleaning Agents, Ozone and Toxic Air Contaminants.
<http://www.arb.ca.gov/research/abstracts/01-336.htm#Disclaimer>

Page 51: “The professional cleaning industry estimates that 85% of the dirt we bring into our homes enters via our shoes or our pets’ paws” US EPA, 2007. Controlling Pollutants and Sources. <http://www.epa.gov/iaq/schooldesign/controlling.html>

Page 52: “According to the report fittingly referred to as the Door Mat Study, lead-contaminated soil from the outside causes almost all the lead dust inside homes. The study also notes that wiping shoes on a mat and removing them at the door cuts lead dust by an impressive 60%.” Small Property Owners of America, n.d.
<http://www.spoa.com/pages/lead3.html>

Page 52: “Synthetic carpets contain chemical stain repellents and mildew treatments, not to mention the toxic chemicals found in the glue and rubber cushioning, which can release noxious fumes for months.” Environmental Science & Technology, 2000. New Indoor Carpet as an Adsorptive Reservoir for Volatile Organic Compounds.
<http://pubs.acs.org/cgi-bin/abstract.cgi/esthag/2000/34/i19/abs/es9910412.html>

Page 55: “The average household does 400 loads of laundry a year” Energy Star, 2007. Clothes Washers: 2007 Partner Resource Guide. http://www.energystar.gov/ia/partners/manuf_res/downloads/2007ClothesWasher_prg.pdf

Page 56: “But chlorine, bleach’s active ingredient, is a powerful disinfectant that can be harmful if inhaled,” US Agency for Toxic Substances & Disease Registry, 2007. ToxFAQs for Chlorine. <http://www.atsdr.cdc.gov/tfacts172.html>

Page 56: “Ninety percent of the energy used for washing clothes goes to heating the water.” US Department of Energy, 2006. Laundry. <http://www1.eere.energy.gov/consumer/tips/laundry.html>

Page 56: “Switching your temperature setting from hot to warm can cut a load’s energy use in half.” US Department of Energy, 2006. Laundry. <http://www1.eere.energy.gov/consumer/tips/laundry.html>

Page 57: “perc” (perchloroethylene, also known as PCE), the highly toxic dry-cleaning fluid. Amazingly, despite the EPA’s listing perc as a hazardous air pollutant in 1997, and a known carcinogen, four of five dry cleaners still use it.” US EPA, 2007. Compliance Assistance for Dry Cleaners. <http://www.epa.gov/region02/capp/dryclean.htm>

Chapter 3 – Clean Plate: Choosing, Eating and Storing Healthier Food

Page 64: “organic product sales are increasing by more than 20% each year.” Organic Trade Association, 2004. Organic Food Facts. <http://www.ota.com/organic/mt/food.html>

Page 67: “While there’s little data on the effect of eating conventionally grown foods, the potential impact of pesticides on the developing brain and nervous system is not a small worry.” National Research Council, 1993. Pesticides in the Diets of Infants and Children. <http://www.nap.edu/openbook.php?isbn=0309048753>

Page 67: “Farmers and others who work with pesticides have elevated rates of asthma, leukemia, and prostate cancer.”

American Journal of Respiratory and Critical Care Medicine, 2002. Pesticide Exposure and Asthma. <http://ajrccm.atsjournals.org/cgi/content/full/165/5/563>

Natural Resources Defense Council, 2000. Trouble on the Farm: Growing Up with Pesticides in Agricultural Communities. <http://www.nrdc.org/health/kids/farm/chap1.asp>

American Journal of Epidemiology, 2003. Use of Agricultural Pesticides and Prostate Cancer Risk in the Agricultural Health Study Cohort. <http://aje.oxfordjournals.org/cgi/reprint/157/9/800.pdf>

Page 67: “And while we’ve made strides in ridding our food of some of the worst chemicals (like DDT).” US Agency for Toxic Substances and Disease Registry, 2002. ToxFAQs for DDT, DDE, and DDD. <http://www.atsdr.cdc.gov/tfacts35.html>

Page 67: “One half of a child’s lifetime exposure to pesticide occurs in the first five years of his or her life.” US EPA, 1999. Pesticides and Child Safety. <http://www.epa.gov/reg3esd1/childhealth/ch4.pdf>

Page 68: “23 – Number of elementary-school children whose urine revealed the presence of pesticides at the beginning of a study in Seattle, Washington. 1 – Number of those same children whose urine contained any detectable traces of pesticides after four days of eating organic food.” Environmental Health Perspectives, 2005. Organic Diets Significantly Lower Children’s Dietary Exposure to Organophosphorus Pesticides. <http://www.ehponline.org/members/2005/8418/8418.html>

Page 68: “That will right away lower your family's pesticide exposure by up to 90%, according to the Environmental Working Group, which did the food analysis.” Environmental Working Group, 2006. Shopper’s Guide: What’s the Difference? <http://www.foodnews.org/methodology.php>

Page 69: “Some types of organic produce have been found to have higher levels of vitamin C and other protective antioxidants.” Proceedings of the Nutrition Society, 2002. Nutritional Quality of Organic Food: Shades of Grey or Shades of Green? <http://www.ingentaconnect.com/content/cabi/pns/2002/00000061/00000001/art00005>

Journal of the Science of Food and Agriculture, 2001. Organic Agriculture: Does It Enhance or Reduce the Nutritional Value of Plant Foods? <http://www3.interscience.wiley.com/cgi-bin/abstract/84002299/ABSTRACT?CRETRY=1&SRETRY=0>

Page 70: “From 1993 to 1997, for example, 50 mid-sized family farms, on average, went under *every day*.” U.S. Department of Agriculture. A National Agricultural Statistics Service, Farm and Land in Farms: Final Estimates, 1993-1997.

Page 71: “The average American family of four tosses out about 1,500 plastic bags a year; each one takes up to 1000 years to decompose.” Center for a New American Dream, 2007. Annapolis Measure Would Ban Plastic Grocery Bags. <http://www.newdream.org/about/mediaclips/wp070207.php>

Page 72: “But you’ve no doubt heard that eating seafood has its risks, due to the buildup of harmful levels of industrial chemicals from pollution and fish farming.” US EPA, 2007. National Lake Fish Tissue Study. <http://epa.gov/waterscience/fish/study/>

Food and Chemical Toxicology, 2002. Organochlorine Chemicals in Seafood: Occurrence and Health Concerns.

<http://www.ncbi.nlm.nih.gov/sites/entrez?db=pubmed&uid=11983271&cmd=showdetailview&indexed=google>

Science, 2004. Global Assessment of Organic Contaminants in Farmed Salmon.
<http://www.sciencemag.org/cgi/content/abstract/303/5655/226>

Page 72: “Two common pollutants—mercury and PCB—are linked to learning and memory problems in children, heart problems, and possibly cancer.” Environmental Health Perspectives, 2005. Public Health and Economic Consequences of Methyl Mercury Toxicity to the Developing Brain.
<http://www.ehponline.org/members/2005/7743/7743.pdf>

US Agency for Toxic Substances and Disease Registry, 2001. ToxFAQs for PCBs.
<http://www.atsdr.cdc.gov/tfacts17.html#bookmark05>

Page 72: “Both mercury and PCBs linger in the body and build up over time: PCBs are removed from the body very slowly (it takes about ten years to get rid of half of last night’s dinner);” Environmental Research, 1992. Changes in PCB serum concentrations among capacitor manufacturing workers.
<http://www.ncbi.nlm.nih.gov/sites/entrez?db=pubmed&uid=1425510&cmd=showdetailview&indexed=google>

Page 72: “for mercury it’s 70 days (so a woman who hopes to become pregnant can clear her body in about one year).” Archives of Environmental Health, 1974. Variations of Biological Half-life in Man.

Page 73: “The most recent government data show that, on average, canned white (albacore) tuna has three times more mercury than canned chunk light tuna, sometimes exceeding the government specified limit.” Environmental Research, 2004. Mercury in Canned Tuna: white Versus Light and Temporal Variation.
http://www.sciencedirect.com/science?_ob=ArticleURL&_udi=B6WDS-4BMC7YR-1&user=10&rdoc=1&fmt=&orig=search&sort=d&view=c&acct=C000050221&version=1&urlVersion=0&userid=10&md5=e2a70196bd9a32d53c8c51be22bbe242

Page 73: “Farmed fish are raised in close quarters and are treated with pesticides and fed antibiotics to minimize lice and disease, a practice that pollutes waterways. Salmon, the third most popular fish in the U.S., is particularly problematic when raised this way. Farmed salmon have been shown to have higher concentrations of pollutants—up to 10 times more PCBs—than wild.” Coastal Alliance for Aquaculture Reform, 2004. Farmed and Dangerous: What Seafood Lovers Should Know About the Salmon They Are Eating.
<http://livingoceans.org/documents/FADReport.04.pdf>

Science, 2004. Global assessment of organic contaminants in farmed salmon.
<http://www.sciencemag.org/cgi/content/abstract/303/5655/226>

Chemosphere, 2002. Preliminary examination of contaminant loadings in farmed salmon, wild salmon and commercial salmon feed.

<http://www.ncbi.nlm.nih.gov/sites/entrez?db=pubmed&uid=11999769&cmd=showdetailview&indexed=google>

Page 78: “For instance, *butylated hydroxyanisole (BHA)*, which keeps oils from going rancid, is listed as a probable human carcinogen by the Department of Health and Human Services, even though the FDA says it’s okay to use.” U.S. Department of Health and Human Services, 2005. Report on Carcinogens, Eleventh Edition.

<http://ntp.niehs.nih.gov/ntp/roc/toc11.html>

Page 78: “The other biggie in this category to look for on labels—and avoid—is *BHT*, or *butylated hydroxytoluene*.” Toxicology, 1981. Enhancement of Tumor Formation in Mouse Lung by Dietary Butylated Hydroxytoluene.

<http://www.ncbi.nlm.nih.gov/sites/entrez?db=pubmed&uid=7281206&cmd=showdetailview&indexed=google>

Page 78: “*Aspartame* (known as NutraSweet and Equal), found in diet sodas and in some kids’ products like vitamins and toothpaste, and *saccharine* (Sweet n’ Low) have a history of controversy. Though no negative health effects have been proven definitively in humans, many scientists have suggested a correlation between these faux-sugars and effects from headaches to cancer.” Environmental Health Perspectives, 2006. First Experimental Demonstration of the Multipotential Carcinogenic Effects of Aspartame Administered in the Feed to Sprague-Dawley Rats.

<http://www.ehponline.org/docs/2005/8711/abstract.html>

US Food and Drug Administration, 1999. Sugar Substitutes: Americans Opt for Sweetness and Lite. http://www.fda.gov/FDAC/features/1999/699_sugar.html

Page 78: “Animal studies have shown a link between high fructose corn syrup and diabetes and high cholesterol, though it’s unclear whether the same is true in humans.” Nutrition Today, 2006. High Fructose Corn Syrups, Part 2: Health Effects.

Page 79: “known also as *partially hydrogenated oils*—are artificial fats that make oil more solid and thereby increase product shelf life. They’re the most harmful fat in the food supply, contributing to heart disease and high cholesterol levels.” New England Journal of Medicine, 2006. Trans Fatty Acids and Cardiovascular Disease.

New England Journal of Medicine, 1999. Trans fatty acids and coronary heart disease.

Page 80: “In particular, avoid FD&C Blues No. 1 and 2; Green No. 3; and Yellow No. 5 & 6.” Townsend Letter, 2007. Artificial Food and Cosmetic Coloring: Hidden Sources of Toxic Metals. <http://www.townsendletter.com/Nov2007/artfood1107.htm>

Page 80: “Sodium nitrites, used to preserve cured and processed meats—bacon, sausage, ham—and keep them looking red and juicy, have been linked to various types of cancer in lab tests.” Journal of the National Cancer Institute, 2005. Meat and Fat Intake as Risk

Factors for Pancreatic Cancer: The Multiethnic Cohort Study.
<http://jnci.oxfordjournals.org/cgi/content/abstract/97/19/1458>

Page 80: “Studies have shown that some people are sensitive to MSG, with reactions ranging from headache and shortness of breath to changes in heart rate (these occur because MSG directly attacks neurons, and at high doses can even kill nerve cells).” US Department of Health and Human Services, 1995. FDA and Monosodium Glutamate (MSG). <http://www.cfsan.fda.gov/~lrd/msg.html>

Endocrinology, 1984. Increased responsiveness of the hypothalamic-pituitary axis after neurotoxin-induced hypothalamic denervation.
<http://www.ncbi.nlm.nih.gov/sites/entrez?db=pubmed&uid=6145582&cmd=showdetailview&indexed=google>

Page 81: “You want to minimize their caffeine exposure for two main reasons: 1) it's a stimulant and kids are energetic enough without it;” Nutrition Bulletin, 2003. The Health Effects of Stimulant Drinks. <http://www.blackwell-synergy.com/doi/abs/10.1046/j.1467-3010.2003.00345.x>

Page 81: “and 2) it's mildly addictive, so you don't want your child experiencing the groaning lethargy of withdrawal.” Journal of the American Academy of Child & Adolescent Psychiatry, 1998. Caffeine Withdrawal in Normal School-Age Children. <http://www.jaacap.com/pt/re/jaacap/abstract.00004583-199808000-00016.htm;jsessionid=HqQL9dKftvx88T2hvsMJ7LMKvgRxV2p64tXhS0vDGMnsKlhXFBY2!1219373867!181195629!8091!-1>

Page 83: “We don't have an exact figure for how pervasive GMO processed foods are, but roughly 60% of processed food in US grocery stores contain at least one soybean product, and more than half of our soybean crops are genetically modified.” Agricultural and Resource Economics Review, 2006. US Consumers' Willingness to Pay for Food Labeled 'Genetically Modified.'
<http://ageconsearch.umn.edu/bitstream/123456789/26332/1/35020299.pdf>

Page 83: “Some research shows worrisome impacts on plants and wildlife, and that food allergens may be transmitted through bioengineering, according to the watchdog group Beyond Pesticides.” Beyond Pesticides, 2007. 10 Reasons to Say No to Genetically Engineered Crops and Foods.
<http://www.beyondpesticides.org/gmos/reportsandpublications/gmofactsheet.htm>

Page 85: “Extra hormones in a child's diet may affect growth and interfere with natural maturation patterns.” Breast Cancer Fund, 2007. The Falling Age of Puberty in US Girls: What We Know, What We Need to Know.
<http://www.breastcancerfund.org/atf/cf/%7BDE68F7B2-5F6A-4B57-9794-AFE5D27A3CFF%7D/The%20Falling%20Age%20of%20Puberty%20report.pdf>

Food and Water Watch, 2007. Say No to rBGH! Keep Artificial Hormones Out of Dairy Products. <http://www.sustainabletable.org/issues/docs/SayNoToRBGH-fww07.pdf>

Page 86: “Ninety percent of American feedlot cattle is injected with growth hormones” Campaign for Food Safety, 1999. “Beef Hormones, Irradiation & Mad Deer: America's Food Safety Crisis Continues.” <http://www.global-reality.com/biotech/articles/news041.4.htm>

Page 86: “Traces of dioxin concentrate in animal fat (including farm-raised fish).” Journal of Animal Science, 1995. A Review of the Significance of Animal Food Products as Potential Pathways of Human Exposures to Dioxins. <http://www.animal-science.org/cgi/content/abstract/73/6/1639>

Page 86: “Charred meat can contain polycyclic aromatic hydrocarbons (PAHs) and heterocyclic amines (HCAs), which can increase the risk of some cancers, according to the National Cancer Institute.” National Cancer Institute, 2004. Heterocyclic Amines in Cooked Meats. <http://www.cancer.gov/cancertopics/factsheet/Risk/heterocyclic-amines>

Journal of Nutrition, 2004. Development of a Food Database of Nitrosamines, Heterocyclic Amines, and Polycyclic Aromatic Hydrocarbons. <http://jn.nutrition.org/cgi/content/full/134/8/2011>

Page 87: “Since its approval by the FDA in 1994, rBGH has been used to boost cows’ milk output by as much as 25%.” E Magazine, 2004. Brave New World: Genetic Engineering, rBGH and Mad Cow Disease. <http://www.emagazine.com/view/?202>

Page 87: “The drug is injected into anywhere from 5 to 30% of cows in the US.” Cornell University Program on Breast Cancer and Environmental Risk Factors, 2000. Consumer Concerns About Hormones in Food. <http://envirocancer.cornell.edu/factsheet/diet/fs37.hormones.pdf>

Page 87: “Use of rBGH has been linked to breast and prostate cancers in animals and the growth factors it produces can be absorbed into the bloodstream.” Journal of the National Cancer Institute, 2001. Re: Role of the Insulin-Like Growth Factors in Cancer Development and Progression. <http://jnci.oxfordjournals.org/cgi/content/full/93/3/238>

Page 87: “a recent survey found that eggs from free range hens had three times the vitamin E, two times the omega-3 fatty acids, and seven times the beta carotene (a form of vitamin A) of non-free range eggs, while averaging one third the cholesterol.” MSNBC, 2007. Pecking Order: Which Eggs are the Best? <http://www.msnbc.msn.com/id/21675726/>

Page 89: “Approximately 12 million Americans now suffer from food allergies, including anywhere from 5 to 7.5% of children;” US Centers for Disease Control, 2005. Allergic

conditions among children: US, 2000-2005.

<http://209.217.72.34/HDAA/TableViewer/tableView.aspx?ReportId=205>.

Page 89: “between 1997 and 2002 the number of children under 5 with peanut allergies doubled, according to the CDC.” Journal of Allergy and Clinical Immunology, 2003. Prevalence of peanut and tree nut allergy in the United States determined by means of a random digit dial telephone survey: A 5-year follow-up study.

Page 89: “What’s causing this allergy increase isn’t well understood. However, one prevailing theory, known as the “hygiene hypothesis,” postulates that we’re *too* clean, thanks to cleansers like antibacterial soaps, which leave the body with few “real” invaders (like viruses and bacteria) to attack, thus the body instead overreacts to very low threats, like nut proteins.” Immunology, 2004. The Increased Prevalence of Allergy and the Hygiene Hypothesis: Missing Immune Deviation, Reduced Immune Suppression, or Both? <http://www.blackwell-synergy.com/doi/abs/10.1111/j.1365-2567.2004.01925.x>

Page 90: “3.1 million Number of school-age children with food allergies”

The Food Allergy and Anaphylaxis Network, 2007. Food Allergy Basics.

<http://www.foodallergy.org/media/FoodAllergyBasics.pdf>

Page 95: “PFOA, the chemical used to make Teflon and other non-stick surfaces, is now found in the blood of nearly every American” US Centers for Disease Control, 2007. National Report on Human Exposure to Environmental Chemicals: Perfluorinated Compounds. http://www.cdc.gov/exposurereport/perfluorinated_compounds2.htm

Page 95: “and has been listed by the EPA as a “likely human carcinogen.” US EPA, 2006. Science Advisory Board Review of EPA’s Draft Risk Assessment of Potential Human Health Effects Associated with PFOA and Its Salts. [http://yosemite.epa.gov/sab%5Csabproduct.nsf/A3C83648E77252828525717F004B9099/\\$File/sab_06_006.pdf](http://yosemite.epa.gov/sab%5Csabproduct.nsf/A3C83648E77252828525717F004B9099/$File/sab_06_006.pdf)

Page 95: “When nonstick pans reach very high temperatures—and tests prove they do—the coating breaks apart and emits toxic particles and gases that may be cancerous, are certainly pollutants, and which are linked to eye and respiratory inflammation (it’s even been blamed for the death of pet birds!).” Environmental Working Group, 2007. Coated Pots and Pans Can Present Health Hazards. <http://www.ewg.org/node/22396>

Page 96: “The lining of microwave popcorn bags are a source of PFOA, the same probable carcinogen found in Teflon. In tests, the FDA found that significant amounts of chemicals leach from the bags to the oil in the popcorn. Though the amounts from any single serving are small, PFOA remains in the body for years and levels can build up.” Food Additives and Contaminants, 2005. Perfluorochemicals: Potential Sources of and Migration from Food Packaging. <http://www.informaworld.com/smpp/content?content=10.1080/02652030500183474>

Page 98: “Many aluminum cans, including liquid formula, are lined with polycarbonate plastic, which contains BPA.” Journal of Health and Science, 2002. Release of Bisphenol A from Food Can Lining Upon Heating. <http://sciencelinks.jp/j-east/article/200218/000020021802A0653963.php>

Page 98: “Americans throw out more than 30 million plastic bottles per day;” Clean Air Council, 2006. Waste Facts and Figures. <http://www.cleanair.org/Waste/wasteFacts.html>

Page 98: “most of these the ubiquitous PET #1 bottles which, while likely posing no immediate health risk, degrade when heated, leaching harmful chemicals” Environmental Science and Technology, 2007. Contamination of Bottled Waters with Antimony Leaching from Polyethylene Terephthalate (PET) Increase upon Storage. <http://pubs.acs.org/cgi-bin/abstract.cgi/esthag/2007/41/i05/abs/es061511+.html>

Page 99: “And baby bottles? In 2007, the advocacy group Environment California released their ‘Toxic Baby Bottles’ report that demonstrated how, when heated, five of the most popular brands made of polycarbonate—the clear, shatterproof plastic—leached BPA at levels found to harm lab animals.” Environment California, 2007. Toxic Baby Bottles: Scientific Study Finds Leaching Chemicals in Clear Plastic Baby Bottles. http://www.environmentcalifornia.org/uploads/Ve/AQ/VeAQsr6MMu4xA3-2ibnr_g/Toxic-Baby-Bottles.pdf

Chapter 4 – The Beauty Part: Natural Body Care

Page 102: “According to the Environmental Working Group, the average person’s grooming routine brings them into contact with 126 different chemicals each day.” Environmental Working Group, 2004. Exposures Add Up – Survey Results. <http://www.cosmeticsdatabase.com/research/exposures.php>

Page 102: “Also the skin is very porous, absorbing up to 60% of what’s put on it” Kat James, 2003. The Truth About Beauty.

Page 107: “*phthalates*, which have been proven to interfere with hormonal and reproductive function in animals. In a 2002 report by the EWG, phthalates were found in nearly three-quarters of the 72 personal care products tested. The number of brands that listed the ingredient on their label? Zero.” Environmental Working Group, 2002. Not Too Pretty: Phthalates, Beauty Products & the FDA.

Page 106: “100, percentage of 289 people, including children and women of childbearing age, who tested positive for more than the safe level of phthalates as determined by the government.” US Centers for Disease Control, 2000. National Report on Human Exposure to Environmental Chemicals. <http://www.cdc.gov/exposurereport/>

Page 107: **1,100+**
the

Number of chemical ingredients banned from cosmetics in

Page 108: “While SLS and SLES are considered safe by the FDA, the big worry is that they can interact with other product ingredients to form carcinogens.” Environmental Working Group, 2008.

<http://www.cosmeticdatabase.com/ingredient.php?ingred06=706110¬hanks=1>

Page 108: “Each year, roughly 6 million elementary school kids are beset with head lice, or pediculosis.” The American Journal of Managed Care, 2004. Head Lice Treatments and the Impact on Managed Care. <http://www.ajmc.com/Article.cfm?Menu=1&ID=2706>

Page 108: “The conventional way to treat head lice is with the prescription shampoo Kwell, a powerful, nerve-attacking insecticide that has even been reported to cause seizures in some children, “ Prescire International, 2006. Lindane: Serious Neurological Effects.

http://www.ncbi.nlm.nih.gov/sites/entrez?Db=pubmed&Cmd=ShowDetailView&TermToSearch=16604739&ordinalpos=1&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVAbstractPlus

Page 110: “Antibacterial soaps are *totally unnecessary*. In fact, they’re a terrible idea: They’re filled with harmful antimicrobial pesticides such as *triclosan*.” Beyond Pesticides, 2004. The Ubiquitous Triclosan: A Common Antibacterial Agent Exposed. <http://www.beyondpesticides.org/pesticides/factsheets/Triclosan%20cited.pdf>

Page 110: “Research shows that washing your hands with regular soap and water work perfectly well for loosening dirt and germs.” Clinical Infectious Diseases, 2007. Consumer Antibacterial Soaps: Effective or Just Risky?

<http://deepblue.lib.umich.edu/dspace/bitstream/2027.42/55477/1/Consumer%20antibacterial%20soaps%20Effective%20or%20just%20risky.web%202007%20aiello.pdf>

Page 110: “According to the “hygiene hypothesis,” the recent leaps in asthma and eczema rates can be partly explained by our cultural insistence on cleanliness and germ-eradication.” Science Daily, 2004. Bacterial Toxin May Protect Infants from Asthma; New Findings Support the Controversial ‘Hygiene Hypothesis.’

<http://www.sciencedaily.com/releases/2004/07/040706075843.htm>

Page 110: “(One recent study by England’s Bristol University investigated how frequently 11,000 toddlers were bathed, showered, or had their hands washed every day and found that “increased levels of hygiene” were linked to higher rates of asthma and eczema for kids from two and four.)” Archives of Disease in Childhood, 2002. Hygiene Levels in a Contemporary Population Cohort are Associated with Wheezing and Atopic Eczema in Preschool Infants. <http://adc.bmj.com/cgi/content/abstract/87/1/26>

Page 110: “Commercial bubble bath is often the cause of “bubble bath cystitis,” a painful irritation of a child’s urethra.” Archives of Disease in Childhood, 2006. Should Bubble Baths be Avoided in Children with Urinary Tract Infections?

<http://www.bestbets.org/cgi-bin/bets.pl?record=01304>

Page 111: “In one study at the Brunel University in England, researchers linked parabens to the possibility that male babies exposed in utero could have lower sperm counts. Another recent British study found traces of parabens in the breast cancer cells of women.” Cornell University Program on Breast Cancer and Environmental Risk Factors, 2004. Parabens: Evidence of Estrogenicity and Endocrine Disruption.

<http://envirocancer.cornell.edu/Bibliography/General/bib.parabens.cfm>

Page 112: “*petroleum distillates*, possible human carcinogens” Environmental Working Group’s Skin Deep Cosmetic Safety Database, 2007. Petroleum Distillates.

http://www.cosmeticsdatabase.com/ingredient.php?ingred06=704787&refurl=%2Fproduct.php%3Fprod_id%3D9068%26

Page 113: “talc can irritate the lungs when inhaled” British Medical Journal, 1991. Inhalation of Baby Powder: An Unappreciated Hazard.

<http://www.pubmedcentral.nih.gov/pagerender.fcgi?artid=1669894&pageindex=1>

Page 113: “and there is evidence that talc may be contaminated with traces of illness-causing asbestos.” US Agency for Toxic Substances and Disease Registry, 2001.

ToxFAQs for Asbestos. <http://www.atsdr.cdc.gov/tfacts61.html>

Page 113: “Conventional wipes are often saturated with alcohol, fragrance, and other skin irritants.”

Environmental Working Group’s Skin Deep Cosmetic Safety Database, 2007. Ingredients to Avoid: Safety Guide to Children’s Personal Care Products.

<http://www.cosmeticsdatabase.com/special/parentsguide/ingredients.php?nothanks=1>

Page 113: “The most common cause of eczema rashes in babies is food allergies, in their diet or that of their nursing mother. This itchy skin inflammation may also be triggered by irritation from harsh soaps, detergents, or synthetic finishes on clothing.” Medscape Dermatology, 2005. Introduction to Atopic Dermatitis.

http://www.medscape.com/viewarticle/519732_2

Page 114: “Skin cancer is the most common cancer in the US;” US Centers for Disease Control, 2007. Skin Cancer, <http://www.cdc.gov/cancer/skin/>

Page 114: “An estimated 40 to 50% of Americans who live to age 65 will get it at least once.” National Institutes of Health, National Human Genome Research Institute, 2007. Learning About Skin Cancer. <http://www.genome.gov/10000184>

Page 114: “The Environmental Working Group recently published an investigation into 831 name-brand sunscreens and discovered that many are neither safe nor effective,

including this doozy: One in eight high-SPF sunscreens does not protect from UVA radiation, the rays linked to skin damage, aging, and potentially skin cancer. Furthermore, many sunscreens contain phthalates, parabens, and other estrogen-like compounds” Environmental Working Group, 2007. Sunscreen Summary – What Works and What’s Safe. <http://www.cosmeticsdatabase.com/special/sunscreens/summary.php>

Page 115: “Skip tanning salons—the UV radiation of a tanning bed is up to 15 times stronger than the sun’s” Medical News Today, 2006. Innovative Public Service Advertisement Campaign Sends Strong Message to Teens About Dangers of Indoor Tanning. <http://www.medicalnewstoday.com/articles/54504.php>

Page 115: “the worry is that nanos are small enough not just to pass through skin but also to penetrate body tissue and organs, potentially causing harm.” Multinational Monitor, 2004. Welcome to NanoWorld: Nanotechnology and the Precautionary Principle Imperative. <http://multinationalmonitor.org/mm2004/09012004/september04corp2.html>

Page 116: “one recent University of Southern California study found that women using permanent hair dye at least once a month for a year or longer more than double their risk of bladder cancer; “ International Journal of Cancer, 2000. Use of Permanent Hair Dyes and Bladder Cancer Risk. <http://www3.interscience.wiley.com/cgi-bin/abstract/76505484/ABSTRACT>

Page 116: “another, from Yale, looking at long-time use, found that permanent hair dyes seemed to increase risk for non-Hodgkins lymphoma.” American Journal of Epidemiology, 2004. Hair Coloring Product Use and Risk of Non-Hodgkin’s Lymphoma: A Population-based Case-Control Study in Connecticut. http://aje.oxfordjournals.org/cgi/content/abstract/159/2/148?maxtoshow=&HITS=10&hits=10&RESULTFORMAT=&title=hair+dye&andorexacttitle=or&andorexacttitleabs=and&andorexactfulltext=and&searchid=1075291054808_352&stored_search=&FIRSTINDEX=0&sortspec=relevance&journalcode=amjepid

Page 116: “Permanent *coal-tar dyes* found in 71 common hair-dye products tested by the EWG have been linked to some types of cancers (coal tar is also found in some dandruff shampoos).” Environmental Working Group, 2005. Use With Discretion. <http://www.ewg.org/node/17847>

Page 116: “11 - Percentage of the 10,000+ chemical ingredients found in cosmetics that have been assessed for health and safety by the Cosmetic Ingredient Review, an industry-funded panel of scientists.”

Environmental Working Group, 2007. Hundreds of Personal Care Products Contain Poorly Studied Nanoscale Materials. <http://www.ewg.org/node/8796>

Page 122: “And fluoride is very effective at that task, cutting down cavities by as much as half.” US Centers for Disease Control, 2007. Oral Health: Preventing Cavities, Gum Disease, and Tooth Loss. <http://www.cdc.gov/nccdphp/publications/aag/oh.htm>

Chapter 5 – Child’s Play: Healthier Toys, Gear, and Clothing

Page 124: “PVC (polyvinyl chloride) plastic, releases toxic fumes into the air”
Jaakkola JJ, Verkasalo PK, Jaakkola N. 2000. Plastic wall materials in the home and respiratory health in young children. *Am J Public Health* 90:797-799.

Jaakkola JJK, Oie L, Nafstad P, Botten G, Samuelsen SO, Magnus P. 1999. Interior surface materials in the home and the development of bronchial obstruction in young children in Oslo, Norway. *Am J Public Health* 89:188-192.

Ponsonby AL, Dwyer T, Kemp A, Cochrane J, Couper D, Carmichael A. 2003. Synthetic bedding and wheeze in childhood. *Epidemiology* 14:37-44.

Page 124: “Studies have proven that phthalates can be hormonally disruptive in animals, and can easily leach out when kids suck or chew on a toy...” *Pediatrics*, 2003. Pediatric Exposure and Potential Toxicity of Phthalate Plasticizers.

<http://pediatrics.aappublications.org/cgi/content/full/111/6/1467>

Page 124: “Another reason to be wary of PVC is that toys made of it may contain lead, which is sometimes used as a stabilizer in this kind of plastic, and can be picked up by children on their hands or in their mouths.” *Current Science*, 2007. Lead and Cadmium in Soft Plastic Toys. <http://www.ias.ac.in/currsci/sep252007/818.pdf>

Page 125: “Opt for the clear silicone pacifiers over the yellow rubber ones, which may contain harmful chemicals.” US Agency for Toxic Substances and Disease Registry, 1989. Public Health Statement for n-Nitrosodimethylamine.

<http://www.atsdr.cdc.gov/toxprofiles/phs141.html>

Page 126: “A recent national study of PVC packaging—that clear plastic casing enveloping toys and electronics, which requires either a Ph.D. or a machete to open—found that more than 60% of samples contained toxic heavy metals, violating legal limits in 19 states.” *Toxics in Packaging Clearinghouse*, 2007. An Assessment of Heavy Metals in Packaging: Screening Results Using a Portable X-Ray Fluorescence Analyzer.

<http://www.toxicsinpackaging.org/>

Page 132: “Also, cotton consumes almost a quarter of the world’s use of insecticides and 10% of herbicides.” United Nations Environment Program, 2007. Textiles: Fashion that Doesn’t Cost the Earth.

http://www.unep.fr/pc/sustain/reports/SCP_Resource_Kit/PDF_FINAL_uk/textiles.pdf

Page 133: “A baby goes through an average of 5,000 to 8,000 diaper changes.” *The Green Guide*, 2006. Product Report: Diapers.

<http://thegreenguide.com/reports/product.mhtml?id=45>

Page 134: “Mostly non-biodegradable, they leave heaps of plastic in landfills for decades—3.6 million tons per year, according to EPA estimates.” US EPA, 2005. Municipal Solid Waste in the United States: 2005 Facts and Figures.
<http://www.epa.gov/msw/pubs/mswchar05.pdf>

Page 134: “They can also emit gases...Since babies inhale more air per pound of body weight than adults do, and are generally more affected by the toxicity of air pollutants, this is troubling.” Acute Respiratory Effects of Diaper Emissions. Archives of Environmental Health, 1999.

Page 139: “Michigan environmental group, Ecology Center, released a study indicating that unwanted chemicals such as chlorine, bromine and lead could be leaching from car seats. Of the 62 car-seat models they tested, 30% had elevated levels of the chemicals; the two biggest culprits were PVCs and flame retardants.” Ecology Center, 2007. HealthyCar.org: The Consumer Guide to Toxic Chemicals in Cars.
<http://www.healthycar.org/reports.php>

Page 140: “Nearly all baby mattresses today use materials that contain harmful ingredients like flame retardants and formaldehyde...” Organic Trade Association, 2007. The O’Mama Report: Better Bedrooms for Healthier Sleep.
http://www.theorganicreport.com/pages/166_healthier_sleep.cfm

Page 141: “*Seattle Post-Intelligencer* published an investigative study in 2000 that turned up asbestos in Crayola, Prang, and Rose Art crayons...” Seattle Post Intelligencer, 2000. Crayola says its Crayons are Asbestos-Free.
<http://seattlepi.nwsource.com/national/cray26.shtml>

Page 142: “Rubber cement, which you probably played with as a kid, is particularly nasty: It contains hexane, a neurotoxin.” US Agency for Toxic Substances and Disease Registry, 1999. Toxicological Profile for n-Hexane.
www.atsdr.cdc.gov/toxprofiles/tp113.pdf

Page 142: “Polymer clays are labeled nontoxic by ACMI, yet they're made of PVCs softened with phthalates...” The Green Guide, 2006. Product Report: Toys.
<http://thegreenguide.com/reports/product.mhtml?id=65&sec=2>

Page 144: “Most lunchboxes are made of vinyl and, according to a recent study by the consumer watchdog group Center for Environmental Health, many contain undesirable levels of lead.” Center for Environmental Health, 2006. Lead in Children’s Lunch Boxes.
<http://www.cehca.org/lunchboxes.htm>

Page 144: “**3 billion** Average number of batteries Americans buy each year for things like radios, toys, cell phones, watches, laptop computers, and portable power tools.” US EPA, 2007. Municipal Solid Waste: Batteries.
<http://www.epa.gov/garbage/battery.htm>

Page 145: “And batteries pose their own problems: They can leak toxic acid that can get onto little hands; when they end up in landfill, they can leach into the soil and groundwater.” Earth911, 2007. How to Recycle and Properly Dispose of Batteries. <http://earth911.org/blog/2007/07/05/how-to-recycle-and-properly-dispose-of-batteries/>

Chapter 6 – Splendor in the Grass: Healthier Gardens, Yards, and Outdoor Spaces

Page 148: “Every year, over 80 million pounds of pesticides are showered on American lawns;” Environment and Human Health, Inc., 2003. Risks from Lawn Care Pesticides: Including Inadequate Packaging and Labeling. <http://www.ehhi.org/reports/lcpesticides/index.shtml>

Page 148: “we use 10 times more pesticides per acre of lawn than farmers use.” The Green Guide, 2007. Product Report: Lawn Care. <http://thegreenguide.com/reports/product.mhtml?id=71>

Page 148: “50 - % of total lifetime pesticide exposure that occur during the first 5 years of a child’s life.” National Research Council, National Academy of Sciences. 1993. Pesticides in the Diets of Infants and Children. <http://www.nap.edu/openbook.php?isbn=0309048753>

Page 149: “children aged six to 11 have been found to possess higher levels of lawn chemicals in their blood than adults, according to the CDC,” US Centers for Disease Control and Prevention. 2003. Second National Report on Human Exposure to Environmental Chemicals. <http://www.cdc.gov/exposurereport/>

Page 149: “cancer in children (and pets) is elevated in relation to the amounts of pesticides and herbicides used on lawns and gardens, among other places.” Archives of Environmental Contamination and Toxicology, 1993. Family Pesticide Use and Childhood Brain Cancer. <http://www.springerlink.com/content/w77584x556w168k1/>

Cancer, 2000. Pesticide Exposures in Children with non-Hodgkin Lymphoma. <http://www3.interscience.wiley.com/cgi-bin/abstract/76501796/ABSTRACT?CRETRY=1&SRETRY=0>

Page 150: “Lawns are the single most irrigated crop in the US, according to NASA—about 40 million acres in total.” TreeHugger, 2007. How to Green Your Gardening. <http://www.treehugger.com/files/2007/06/how-to-green-your-gardening.php>

Page 158: “Since most pesticides are designed to be toxic to the nervous system of an insect, there is concern that exposure to common lawn and garden pesticides might compromise the human nervous system, and lead to unwanted consequences.” US EPA, 2007. Pesticides: Health and Safety. <http://www.epa.gov/opp00001/health/human.htm>

Page 158: “Other studies suggest that use of household and garden pesticides increases the risk of childhood leukemia.” Environmental Health Perspectives, 2002. Critical Windows of Exposure to Household Pesticides and Risk of Childhood Leukemia.

<http://www.ehponline.org/docs/2002/110p955-960ma/abstract.html>

Page 158: “Pregnant women, in particular, should be especially wary, since exposure may contribute to complications and birth defects.” Environmental Health Perspectives, 2004. Low-Dose Agrochemicals and Lawn-Care Pesticides Induce Developmental Toxicity in Murine Preimplantation Embryos.

<http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=1241965>

Environmental Health Perspectives, 1999. Exposures of Children to Organophosphate Pesticides and Their Potential Adverse Health Effects.

<http://books.google.com/books?hl=en&lr=&id=HD2JJzD3CDAC&oi=fnd&pg=PA409&dq=Environmental+Health+Perspectives,+1999.+Exposures+of+Children+to+Organophosphate+Pesticides+and+Their+Potential+Adverse+Health+Effects.+&ots=M1xnAUbfN4&sig=yF8NFoShWWxjEJ1EzkkHrHpJBjYr8#PPA384,M1>

Page 158: “10 - Number of times indoor levels of pesticide can go up after an outdoor application.” Ginsberg & Toal, 2006. What’s Toxic, What’s Not.

http://www.whatstoxic.com/THE_BOOK.html

Page 159: “Despite the ubiquity of citronella and herbal products, a test published in the *New England Journal of Medicine* demonstrated they kept mosquitoes away for less than 20 minutes.” The New England Journal of Medicine, 2002. Comparative Efficacy of Insect Repellents Against Mosquito Bites.

<http://content.nejm.org/cgi/content/full/347/1/13>

Page 159: “The only natural product whose effectiveness exceeded one hour is a soybean oil product called Bite Blocker for Kids, which the US Department of Agriculture found to be as effective as a 15% concentration of DEET and lasted for 4-8 hours.” US Department of Agriculture, 2004. Laboratory Evaluation of Mosquito Repellents Against *Aedes albopictus*, *Culex nigripalpus*, and *Ochlerotatus triseriatus*.

http://www.homs.com/pdf/July2004_USDA_marked.pdf

Page 159: “The botanical product Repel, with lemon eucalyptus derivatives, was found by the CDC to be as effective as low concentrations of DEET.” US CDC, 2006. Updated Information Regarding Insect Repellents.

<http://www.cdc.gov/ncidod/dvbid/westnile/RepellentUpdates.htm>

Page 164: “The Environmental Working Group has shown that high levels of arsenic leach out of pressure-treated wood for years after installation—which happens when rain pools on it, bringing arsenic to the surface. The EWG also found that the amount of arsenic that testers wiped off an area of wood—about the size of a four-year-old's hand—exceeded what the EPA allows in a glass of water. Just as worrisome, arsenic in the soil from two of every five backyards or parks tested exceeds EPA's Superfund cleanup

levels.” Environmental Working Group, 2007. Health/Toxics: Arsenic in Treated Wood. <http://www.ewg.org/featured/232>

Page 166: “Some play sand used in sandboxes and around swing sets contain asbestos or very fine crystalline silica. The most dangerous play sand is made from crushed quarry rock.” The Safe Sand Company, n.d. Playsand Information. <http://www.safesand.com/information.htm>

The Green Guide, 2007. True Grit: Safer Sand and Play Sets. <http://www.thegreenguide.com/doc/120/play>

Page 167: “In a study on the safety of thousands of playgrounds, the US Public Interest Research Group and Consumer Federation of America found that half of the playgrounds they tested had peeling, chipping or cracking paint on equipment surfaces.” US PIRG Education Fund, 2002. Playing It Safe: The Sixth Nationwide Safety Survey of Public Playgrounds. <http://www.consumerfed.org/pdfs/PlayingItSafeJune2002.pdf>

Page 168: Chlorine in swimming pools kills bacteria and other microorganisms in water but can be inhaled or absorbed and cause irritated skin and burning eyes.

Page 168: “It can also lead to respiratory problems and an increased risk of asthma.” European Respiratory Journal, 2002. Indoor Swimming Pools, Water Chlorination and Respiratory Health. <http://erj.ersjournals.com/cgi/content/full/19/5/790>

Occupational and Environmental Medicine, 2003. Lung hyperpermeability and asthma prevalence in schoolchildren: unexpected associations with the attendance at indoor chlorinated swimming pools. <http://oem.bmj.com/cgi/content/abstract/60/6/385>

Page 169: “Eighty-five percent of the dirt in homes is brought in from the outside, on our shoe bottoms or the paws of pets.” Capital Health Edmonton Area, 2006. How to Have a Healthy Home. <http://www.capitalhealth.ca/AboutUs/ResourceLibrary/YourHealthMagazine/SeptOct2006/HealthyHome.htm>

Page 169: “Another study has shown that virtually all lead dust inside homes is caused by lead-contaminated soil from outside. Wiping shoes on a mat and removing them at the door cuts lead dust by 60%.” J.W. Roberts, D.E. Camann and T.M. Spittler, presented at the 84th Annual Meeting of the Air and Waste Management Association, June 1999. Reducing Lead Exposure from Remodeling and Soil Track-In in Older Homes. Summarized at <http://www.spoa.com/pages/lead3.html>

Chapter 7 – Every Sip, and Breath, You Take: Clean Water & Air Solutions

Page 171: “and air *indoors*, where we spend 90% of our time, and where the concentration of pollutants in the air is two to five times greater, on average, than it is

outdoors” US EPA, 2007. The Inside Story: A Guide to Indoor Air Quality.
<http://www.epa.gov/iaq/pubs/insidest.html>

Page 172: “Lead can leach into water that sits overnight in plumbing, particularly in older houses.” US EPA, 1993. Actions You Can Take to Reduce Lead in Drinking Water.
<http://www.epa.gov/OGWDW/lead/lead1.html>

Page 172: “Lead is particularly toxic to kids and has been linked to decreased intelligence and developmental disabilities.” US EPA, 2006. Lead in Drinking Water: Basic Information. <http://www.epa.gov/OGWDW/lead/basicinformation.html#healtheffects>

Page 172: “Since many contaminants are more apt to leach into hot tap water than cold, use cold water for cooking, drinking, or—very important—mixing infant formula.” DC Water and Sewer Authority, n.d. Water Quality FAQs.
<http://www.dcwasa.com/waterquality/faqs.cfm>

Page 173: “chemical byproducts called THMs, which are carcinogens.” Dissertaion Abstracts International Part B: Science and Engineering, 1982. Stochastic Prediction of Trihalomethane Production in Chlorinated Drinking Water.
<http://mdl.csa.com/partners/viewrecord.php?requester=gs&collection=ENV&recid=355167&q=Trihalomethane+carcinogenic&uid=791885774&setcookie=yes>

Page 174: “260 – Number of contaminants detected in water, over half of which have no safety standards.” Environmental Working Group, 2005. A National Assessment of Tap Water Quality. <http://www.ewg.org/tapwater/findings.php>

Page 176: “137 million petroleum-based plastic bottles pile up in landfills every day” Container Recycling Institute, 2006. PET Beverage Sales and Wasting.
<http://www.container-recycling.org/images/plastic/graphs/PETwastesale-unit-96-06.gif>

Page 176: “and take at least 1,000 years to decompose,” E Magazine, 2005. Better Bottles. <http://www.emagazine.com/view/?2844>

Page 176: “the Natural Resources Defense Council found that the water from one-third of the 103 bottled water companies tested were actually of *lower* quality than tap” Natural Resources Defense Council, 1999. Bottled Water: Pure Drink or Pure Hype?
<http://www.nrdc.org/water/drinking/bw/bwinx.asp>

Page 176: “Polycarbonate can leach its primary building block—the suspected hormone disruptor, bisphenol-A—into the liquid; the hotter or older it gets, the likelier the leaching.” Environmental Health Perspectives, 2003. Bisphenol A is Released from Used Polycarbonate Animal Cages into Water at Room Temperature.
<http://www.ehponline.org/members/2003/5993/5993.html>

Page 178: “**4:1** approximate ratio of volume of water used for a typical bath to that used for a five-minute shower (using a conventional, non-low-flow shower head).” US EPA, 2007. Use Your WaterSense. <http://www.epa.gov/watersense/water/simple.htm>

Page 179: “Since 1985, rates of childhood asthma have more than doubled.” US Centers for Disease Control, National Center for Health Statistics, 2006. State of Childhood Asthma, United States: 1980-2005. <http://www.cdc.gov/nchs/data/ad/ad381.pdf>

Page 182: “American Academy of Allergy, Asthma & Immunology, allergic rhinitis or hay fever accounts for 2 million lost days of school annually” American Academy of Allergy, Asthma & Immunology, 2007. Allergy Statistics. http://www.aaaai.org/media/resources/media_kit/allergy_statistics.stm

Page 182: “**6,500,000** the number of American children affected by asthma” US Centers for Disease Control, National Center for Health Statistics, 2006. State of Childhood Asthma, United States: 1980-2005. <http://www.cdc.gov/nchs/data/ad/ad381.pdf>

Page 187: “formaldehyde and benzene. In the short term, these chemicals can make your eyes and throat hurt or, in people who are very sensitive, set off an asthma attack. In the long term, they're suspected to cause more serious health effects, including cancer.” US Agency for Toxic Substances and Disease Registry, 1999. ToxFAQs for Formaldehyde. <http://www.atsdr.cdc.gov/tfacts111.html>

US Agency for Toxic Substances and Disease Registry, 2007. ToxFAQs for Benzene. <http://www.atsdr.cdc.gov/tfacts3.html>

Page 188: “1,000 -estimated number of times worse indoor air is than outdoor air, immediately after painting with conventional paint” US EPA, 2007. The Inside Story: A Guide to Indoor Air Quality. <http://www.epa.gov/iaq/pubs/insidest.html>

Page 189: “radon, which is the second-leading cause of lung cancer.” US EPA, 2007. A Citizen’s Guide to Radon: The Guide to Protecting Yourself and Your Family from Radon. <http://www.epa.gov/radon/pubs/citguide.html>

Page 189: “CO, produced when any fuel (e.g., gas, fuel oil, natural gas, kerosene, wood, coal, charcoal) is burned, can accumulate rapidly indoors, and may cause headaches, dizziness, and can even be fatal within minutes.” US EPA, 2007. Basic Information: Carbon Monoxide. <http://www.epa.gov/radon/pubs/citguide.html>

Page 189: “Open the oven door and you get a blast of nitrogen dioxide, a pollutant that can irritate breathing” National Library of Medicine, 2007. Tox Town: Nitrogen Oxides. http://toxtown.nlm.nih.gov/text_version/chemicals.php?id=19

Page 191: “ozone generators have been found to create byproducts that are more toxic” US EPA, 2007. Ozone Generators that are Sold as Air Cleaners.

<http://www.epa.gov/iaq/pubs/ozonegen.html#are%20ozone%20generators%20effective%20in%20controlling%20pollution>

Page 192: “Natural Resources Defense Council study of 14 commonly available air fresheners, hormone-disrupting phthalates were found in 12 of them” Natural Resources Defense Council, 2007. Hidden Hazards in Air Fresheners.

<http://www.nrdc.org/health/home/airfresheners.asp>

Page 193: “Scented candles release more chemicals (such as formaldehyde and benzene) and soot when burned than unscented ones.” Fragranced Products Information Network, 2005. Scented Products as Sources of VOCs: Implications for Susceptible Populations.

<http://www.fpinva.org/AWMA2005.htm>

Page 193: “A study conducted for NASA identified several houseplants that reduce the concentration of common contaminants in the air.” NASA, 1989. Interior Landscape Plants for Indoor Air Pollution Abatement.

http://ntrs.nasa.gov/archive/nasa/casi.ntrs.nasa.gov/19930073077_1993073077.pdf

Center for Environmental Oncology, University of Pittsburgh Cancer Institute, 2007. Greening Indoor Air.

<http://environmentaloncology.org/files/file/Publications/Newsletters/2007-Fall-News.pdf>

Chapter 8 – BFF: Raising a Green Pet

Page 198: “Regular washing can reduce allergens by up to 84%.” Humane Society of Greater Miami, 2005. Coping with Allergies to Pets.

<http://www.humanesocietymiami.org/articles.htm>

Page 200: “Within eight hours after a flea lays eggs on your cat or dog, about 70% of the eggs have fallen onto the carpet or the pet’s bedding” Annual Review of Entomology, 1997. The Biology, Ecology, and Management of the Cat Flea.

<http://arjournals.annualreviews.org/doi/abs/10.1146/annurev.ento.42.1.451>

Page 200: “A University of California study showed that vacuuming catches about 96% of adult fleas.” Journal of Economic Entomology, 1986. Distribution and control of cat fleas in homes in southern California (Siphonaptera: Pulicidae).

Page 200: “A non-manufacturer study found traps effective at capturing 10% of adult fleas;

studies sponsored by trap makers found they’re up to 95% effective.” Journal of Pesticide Reform, 2003. Managing Fleas Without Poison. <http://www.pesticide.org/fleas.html>

Page 203: “Not only is animal waste unsightly and stinky, it transmits diseases and parasites to people. Two intestinal parasites, ascarids and hookworms, are common in puppies and kittens. The CDC determined that eating dirt is the primary way of

contracting these parasites.” The Green Guide, 2005. Product Report: Pet Products.
<http://thegreenguide.com/reports/product.mhtml?id=48&sec=2>

Page 203: “Clay is strip-mined and filled with silica dust that may irritate your cat’s—and child’s—lungs. The clumping agent, sodium bentonite (used also in grouting, sealing, and plugging agents), can swell up to 18 times its dry size, and poison your cat if it eats enough.” E Magazine, 1998. From Rainforests to Kitty Litter.
<http://www.emagazine.com/view/?624>

Page 203: “Toxoplasmosis, a potentially serious disease that can infect the fetus and cause birth defects or miscarriage, is transmitted through cat feces.” The Green Guide, 2005. Product Report: Pet Products.
<http://thegreenguide.com/reports/product.mhtml?id=48&sec=2>

Page 210: “some studies have linked lawn chemicals to bladder cancer in canines.” Journal of the American Veterinary Medical Association, 2004. Herbicide Exposure and the Risk of Transitional Cell Carcinoma of the Urinary Bladder in Scottish Terriers.
<http://avmajournals.avma.org/doi/abs/10.2460/javma.2004.224.1290?cookieSet=1&journalCode=javma>

Page 211: “Chloride salts, the most common de-icer, irritate the skin of pets and can even poison them if they lick their feet. Avoid carbonyl diamide—a.k.a. urea—to melt ice: It releases nitrate when it enters the water supply, causing algae blooms.” The Green Guide, 2004. A Pet-Friendly De-Icer? <http://www.thegreenguide.com/doc/ask/ice>

Page 213: “Puppies may pass the bacterium *Campylobacter* and a variety of other parasites, which can cause diarrhea and rashes or illness in humans, especially children under five.” American Academy of Family Physicians, 2007. Pet-Related Infections.
<http://www.aafp.org/afp/20071101/1314.pdf>

Chapter 9 – Safe House: Healthy Home Improvement

Page 215: “You can also find mattresses made of natural latex, which deters dust mites and bacteria, and is stain- and water-proof.” Debra Lynn Dadd, 2006. Choosing a Natural Bed. <http://www.lulu.com/content/312054>

Page 223: “Standard household paints release solvents, or VOCs (volatile organic compounds), such as benzene and formaldehyde, which can cause breathing problems and may even subtly contribute to other health problems, like damage to the nervous system or cancer.” US Agency for Toxic Substances and Disease Registry, 2007. ToxFAQs for Benzene. <http://www.atsdr.cdc.gov/tfacts3.html>

US Agency for Toxic Substances and Disease Registry, 1999. ToxFAQs for Formaldehyde. <http://www.atsdr.cdc.gov/tfacts111.html>

Page 226: “‘manufactured’ woods—pressed wood and plywood (recognizable by their

layer-cake cross-section), and particleboard and chipboard (which looks like wood chips stuck in a cake). These contain glues that give off formaldehyde, a probable carcinogen.” US EPA, 2007. Indoor Air Quality Basic Information: Formaldehyde.

<http://www.epa.gov/iaq/formalde.html>

Page 232: “All electronics contain toxic chemicals and heavy metals—flame retardants, lead, cadmium, and mercury, to name a few.” GreenPeace, 2006. Toxic Tech: The Dangerous Chemicals in Electronic Products.

<http://www.greenpeace.org/raw/content/international/press/reports/toxic-tech-chemicals-in-elec.pdf>

Page 232: “While exposure likely isn’t a problem for casual users, some elevated levels of flame retardants have shown up in the blood of super-frequent users.” Silicon Valley Toxics Coalition, 2002. Just Say No to E-Waste: Background Document on Hazards and Waste from Computers. <http://svtc.igc.org/cleancc/pubs/sayno.htm>

Page 232: “Electronics are the number one source of individual hazardous waste.” Earth911, 2007. E-Waste: The Final Frontier. <http://earth911.org/blog/2007/10/11/e-waste-the-final-frontier/>

Page 233: “70 – percentage of all hazardous waste that comes from electronics.” Worldchanging.com, 2005. Sustainable Electronics.

<http://www.worldchanging.com/archives/002088.html>

Page 235: “It’s not unusual for home duct systems to leak 25% or more of its air during operation,” US Department of Energy, 1999. Improving the Efficiency of Your Duct System. <http://www.eere.energy.gov/buildings/info/documents/pdfs/27630.pdf>

Page 236: “excessive exposure to magnetic fields is associated with an elevation in childhood leukemia.” Environmental Health Perspectives, 1995. Childhood Leukemia and Residential Exposure to Weak Extremely Low Frequency Magnetic Fields.

<http://www.ehponline.org/members/1995/Suppl-2/feychting-full.html>

Chapter 10 – It’s All Good: How to Grow Your Impact

Page 254: “In 1960, the average American created 2.7 pounds of waste daily; in 2001, it was 4.4 pounds.” US EPA, 2003. EPA’s Draft Report on the Environment 2003.

<http://www.epa.gov/roe/roe/pdf/roeLand.pdf>

Page 256: “(there are over three times as many obese children aged 6 to 11 today than in 1970).” Institute of Medicine, 2004. Childhood Obesity in the United States: Facts and Figures.

<http://www.iom.edu/Object.File/Master/22/606/FINALfactsandfigures2.pdf>

Page 256: “Between 1981 and 1997, the number of hours American kids aged 6 to 8 spent playing outside dropped by four per week, while the number of hours in school

went up by five.” University of Michigan Institute for Social Research, 2000. Changes in American Children’s Time, 1981-1997.

<http://ceel.psc.isr.umich.edu/pubs/papers/ceel013-00.pdf>

Page 256: “American children spend an average of 30 minutes of unstructured time outdoors...*each week.*” University of Michigan Institute for Social Research, 2000. Changes in American Children’s Time, 1981-1997.

<http://ceel.psc.isr.umich.edu/pubs/papers/ceel013-00.pdf>

Page 256: “In a study of 830 mothers, 70% said they played outdoors daily as kids, compared with 31% of their children.” Contemporary Issues in Early Childhood, 2004. An Investigation of the Status of Outdoor Play.

<http://www.wwwords.co.uk/rss/abstract.asp?j=ciec&aid=1634>

Page 256: “Kids 8 to 18 spend six-plus hours a day with TV, electronic games, the computer, and other screens and audio.” Kaiser Family Foundation and the Centers for Disease Control and Prevention (cited in USA Today, July 12, 2005).

Page 256: “more homework (for America’s youngest children, the amount has tripled since 1981” University of Michigan Institute for Social Research, 2000. Changes in American Children’s Time, 1981-1997.

<http://ceel.psc.isr.umich.edu/pubs/papers/ceel013-00.pdf>

Page 257: “In one intergenerational study, the number of people who reported that the outdoors was the most influential environment of their childhood dropped from 96% to 46%.)” Environment and Behavior, 1991. The Landscapes of Childhood — The Reflection of Childhood’s Environment in Adult Memories and in Children’s Attitudes.

<http://eab.sagepub.com/cgi/content/abstract/23/4/395>

Page 257: “82 - % of mothers, in one study, who cite crime and safety concerns as factors for preventing their kids from playing outside.” Contemporary Issues in Early Childhood, 2004. An Investigation of the Status of Outdoor Play.

<http://www.wwwords.co.uk/rss/abstract.asp?j=ciec&aid=1634>

Page 257: “38 - % that violent victimization of children has dropped since 1975.” Duke University, 2005. Child Well Being Index. <http://www.soc.duke.edu/~cwi/>